Application of Antimicrobial Peptides for Disease Control in Plants

Yu-Jin Jung^{1,2}, Kwon-Kyoo Kang^{1,2*}

¹Department of Horticulture, Hankyong National University, Anseong 456-749, Korea

²Institute of Genetic Engineering, Hankyong National University, Anseong, 456-749, Korea

ABSTRACT A large number of antimicrobial peptides (AMPs) from different organisms have been characterized to date. AMPs are small molecular weight proteins with broad spectrum antimicrobial activity against bacteria, viruses, and fungi. Several diseases in plants could result in serious losses and decreasing quality and safety of agricultural products. Being an agent for plant defense, AMPs kill target cells through diverse mechanisms once in a target microbial membrane. Cathelicidins, defensins and thionins are the three major groups of epidermal AMPs in human and plants. Plant AMPs are structurally and functionally diverse and can be directed against other organisms, like herbivorous insects. The biological activity of plant AMPs primarily depends on interactions with membrane lipids. Several antimicrobial peptides have been expressed in transgenic plants to confer disease protection. Antimicrobial peptides are interesting compounds that can be efficiently exploited for disease control in plants in a way that complies with the strict regulations on the efficacy and safety of disease control strategy.

Keywords Antimicrobial peptide, Cathelicidins, Defensins, Thionins, Plant protection

INTRODUCTION

Plant diseases caused by viruses, bacteria and fungi affect crops, and have an effect on significant losses or decrease the quality and safety of agricultural products (Agrios 2005). In order to reduce yield loss, development of novel breeding strategies is urgently needed. Plants have long been known to exhibit mechanisms that enabled them to detect and defend against microbial attack. In response to microbial attack, plants have activated a complex series of responses that lead to the local and systemic induction of a broad-spectrum of antimicrobial defenses (Montesinos 2007).

In human medicine, antimicrobial peptides (AMPs) have been considered as potential therapeutic sources of future antibiotics because of their broad-spectrum activities and different mechanisms of action compared to conventional antibiotics. Although AMPs possess considerable benefits as new generation antibiotics, their clinical and commercial development still have some limitations, such as potential

toxicity, susceptibility to proteases, and high cost of peptide production. In order to overcome those obstacles, extensive efforts have been carried out.

Antimicrobial peptides (AMPs) are endogenous polypeptides produced by multicellular organisms in order to protect a host from pathogenic microbes. AMPs are also defined as host defense peptides because of their essential role in constituting the innate immunity system (Zasloff 2002; Nguyen *et al.* 2011). AMPs are short sequence peptides with generally fewer than 50 amino acid residues, most of which have antimicrobial activity against a broad spectrum of pathogens. In general, when AMPs are folded in membrane mimetic environments, one side of AMPs is positively charged (mainly due to lysine and arginine residues) and the other side contains a considerable proportion of hydrophobic residues (Zasloff 2002; Hancock *et al.* 1997; Hancock and Lehrer 1998). The cationic properties of AMPs are implicated in their selective interaction with the negatively charged surfaces of microbial membranes,

Received March 26, 2014; Revised March 28, 2014; Accepted March 29, 2014; Published March 31, 2014

*Corresponding author Kwon-Kyoo Kang, kykang@hknu.ac.kr, Tel: +82-31-670-5104, Fax: +82-31-670-5109

resulting in the accumulation of AMPs on the membrane surface. And their hydrophobic portions are responsible for the interaction with hydrophobic components of the membrane. From this complex interaction with the membrane, major rearrangements of its structure occur, which may result from the formation of peptide-lipid specific interactions, the peptide translocation across the membrane and interaction with intracellular targets or the most common mechanism, a membranolytic effect (Nguyen *et al.* 2011; Hancock 1997).

AMPs show broad-spectrum antimicrobial activities against various microorganisms, including Gram-positive and Gram-negative bacteria, fungi, and viruses (Zasloff 2002). Most AMPs, with their amphipathic nature, directly act on the membrane of the pathogen. They are a first line of defense in plants and animals, and resistance against them is much less observed compared with current antibiotics (Hancock 1997). AMPs from various sources have been demonstrated to confer resistance against fungal and bacterial pathogens in an array of genetically engineered plant species, including *Arabidopsis* (Maloy 1995), tobacco (Ramamoorthy *et al.* 2006; Kang *et al.* 2009; Park *et al.* 2002), Chinese cabbage (Jung *et al.* 2012), rice (Gordon *et al.* 2005; Oyston *et al.* 2009), tomato (Lupetti *et al.* 2000), cotton (Lupetti *et al.* 2003), potato (Brouwer *et al.* 2011), pear (Hwang *et al.* 1998), banana (Ramamoorthy *et al.* 2006) and hybrid poplar (Nguyen *et al.* 2005).

In this review, we describe the potentials of AMPs as novel disease control agent in plants, and its structural functions that relate to the mechanism of its inhibitory action to invading pathogen.

History and structural information of antimicrobial peptides

Antimicrobial peptides (AMPs) are oligopeptides with a varying number (from five to over a hundred) of amino acids. The discovery of AMPs dates back to 1939, when Dubos (Dubos 1939) extracted an antimicrobial agent from a soil *Bacillus* strain. This extract was demonstrated to protect mice from pneumococci infection. In the following year, Hotchkiss and Dubos (Hotchkiss and Dubos 1940) fractionated this extract and identified an AMP which was named gramicidin. Despite some reported toxicity associated

with intraperitoneal application (Hotchkiss and Dubos 1940), gramicidin was found effective for topical treatment of wounds and ulcers (Van Epps 2006). In 1941, another AMP, tyrocidine, was discovered and found to be effective against both Gram-negative and Gram-positive bacteria (Dubos and Hotchkiss 1941). However, tyrocidine exhibited toxicity to human blood cells (Rammelkamp 1942). In the same year, another AMP was isolated from a plant *Triticumaestivum* (Balls 1942), which was later named purothionin and found effective against fungi and some pathogenic bacteria (Ohtani *et al.* 1977). The first reported animal-derived AMP is defensin, which was isolated from rabbit leukocytes in 1956 (Hirsch 1956). In the following years, bombinin from epithelia and lactoferrin from cow milk were both described (Groves *et al.* 1965). During the same time, it was also proven that human leukocytes contain AMPs in their lysosomes (Zeya and Spitznagel 1963). In total, more than 5,000 AMPs have been discovered or synthesized up to date (Zhao *et al.* 2013). Natural AMPs can be found in both prokaryotes (e.g., bacteria) and eukaryotes (e.g., protozoan, fungi, plants, insects, and animals) (Conlon and Sonnevend 2010). Most AMPs are produced by specific cells at all times, while the production of some AMPs is inducible.

AMPs can be classified into four groups based on their structures: α -helical peptides, β -sheet peptides which contain two or more disulfide bonds, β -hairpin or loop which possess single disulfide bond and/or cyclization of the peptide chain, and extended peptides (Zasloff 2002; Nguyen *et al.* 2011; Hancock and Lehrer 1998). Among these structural groups, α -helix and β -sheet structures are more common and α -helical peptides are the most studied AMPs to date (Arnold 1977) (Table 1). The α -helical AMPs, including magainin, cecropin, and pexiganan, constitute a representative class of AMPs that are the most well established in structure-activity relationships. This group of peptides is usually unstructured in aqueous solution and forms amphipathic helices in membranes or membrane-mimicking environments (Seo *et al.* 2012). Most α -helical AMPs disrupt bacterial membranes, and several mechanisms of action employed by various AMPs have been proposed. The α -helical amphipathic peptides form barrel-like bundles in the bacterial membranes, and

these transmembrane clusters line amphipathic pores (barrel-stave model). Many α -helical AMPs, including some magainins and cecropins, can disrupt bacterial membranes by forming carpet-like clusters of peptides. The peptides adsorb and align in parallel to the surface of bacterial membranes, then the membranes are collapsed into micelle-like structures by high concentrations of

peptides (Fig. 1). The α -helical AMPs, such as some magainins and protegrins, form toroidal pores to disrupt the bacterial membranes (toroidal pore model) (Zasloff 2002; Nguyen *et al.* 2011; Hancock and Lehrer 1998). The β -sheet AMPs, such as α -, β -defensins, and protegrin, are stabilized by disulfide bridges, and form relatively rigid structures. Many of β -sheet AMPs exert their antimicrobial

Table 1. Representative antimicrobial peptides.

Type	AMPs	Size	Characteristic	Origin	Activity	Reference
α -helical peptides of Lack in cysteine	Magainin	~2.5 kDa, 23 amino acid (aa)	Lys rich	analog of frogs	Gram positive/negative bacteria, fungi, parasite	Gesell <i>et al.</i> 1997
	Cecropins	~4 kDa, 30~45 aa	Lys rich, Amidation of C-terminal	Blood of insects	Gram positive/negative bacteria	Holak <i>et al.</i> 1988
	Cecropin P1	~4 kDa, 30~45 aa	Amidation of C-terminal	paneth cell from small intestine of pigs	Gram negative bacteria	Sipos <i>et al.</i> 1992
	Buforin	39 aa	Similarity to C-terminal of Histone IIa	Stomach of American Bull Frog	Gram positive/negative bacteria, fungi,	Yi <i>et al.</i> 1996
	hCAP18/ LL-37	18 kDa, 37 aa	Helical C-terminal	humans	Gram positive/negative bacteria, fungi,	Gudmundsson <i>et al.</i> 1996
Cationic peptide enriched for specific amino acid	Bac5, Bac7	43 or 59 aa	Lack in cysteine residue, rich in proline, arginine, phenylalanine, glycine, tryptophan	Neutrophils of cattle	Gram positive/negative bacteria	Scocchi <i>et al.</i> 1994
	PR39	39 aa		small intestine of pigs		Agerberth <i>et al.</i> 1991
	Indolicidin	13 aa		Neutrophils of cattle		Selsted <i>et al.</i> 1992
β -hairpin or loop due to the presence of a single disulfide bond	Dodecapeptide	12 aa	and/or cyclization of the peptide chain	Neutrophils of cattle	Gram positive/negative bacteria	Romeo <i>et al.</i> 1988
	Brevinins	20~34 aa		amphibians		Conlon <i>et al.</i> 1999
	Raqaalexin	20 aa		amphibians		Clark <i>et al.</i> 1994
β -sheet peptides due to the presence of 2 or more disulfide bonds	Defensins	~4 kDa, 29~45 aa	Arg-rich, salt or iron sensitive activity	Neutrophils of rabbits and humans	Gram positive/negative bacteria, fungi, cytotoxicity	Pardi <i>et al.</i> 1992
	β -Defensin	38~42 aa		Bovine leukocytes		Zimmermann <i>et al.</i> 1995
	Protegrin	16~18 aa	S-S bond	Porcine leukocytes		Fahrner <i>et al.</i> 1996

activities by disrupting bacterial membranes. They are perpendicularly inserted or tilted into the lipid bilayer to form toroidal pores, and hydrophilic regions of the peptides are associated with the polar head groups of the membranes (Nguyen *et al.* 2011). While most AMPs belong to one of the four classes, some AMPs do not belong to any of these groups. Some of which contain two different structural components (Brouwer *et al.* 2011). Also, many peptides form their active structure only when they interact with the membranes of target cells.

These AMP gene families have little in common other than their ability to kill microbes. Also, this structure appears critical to their ability to interact with the negatively charged microbial membrane in the initiation of antimicrobial effects.

The major groups of AMPs: Cathelicidins, Defensins and Thionins

Like AMPs of animal origin, the molecular diversity of AMPs from plants is striking (Padovan *et al.* 2010). The most common classes are thionins, defensins, and lipid transfer proteins. Plant AMPs share the following important features: They are small cationic peptides with molecular masses of 2–10 kDa. The structures of these small peptides are stabilized through formation of 2–6 disulfide bridges. The activities of plant AMPs are primarily directed against fungal, oomycete, and bacterial microorganisms, but certain members of a class can be directed against other targets, including herbivorous insects. Different classes of plant AMPs are as follows.

Cathelicidins form a distinct class of proteins present in the innate immunity of mammals (Zanetti 2004). Similar to

Fig. 1. Schematic representation of some action mechanisms of membrane-active AMPs.

(A) Barrel-Stave model. AMP molecules insert themselves into the membrane perpendicularly. (B) Carpet model. Small areas of the membrane are coated with AMP molecules with hydrophobic sides facing inward leaving pores behind in the membrane. (C) Toroidal pore model. This model resembles the Barrel-stave model, but AMPs are always in contact with phospholipid head groups of the membrane. The blue color represents the hydrophobic portions of AMPs, while the red color represents the hydrophilic parts of the AMPs. (Bahar and Ren 2013).

defensins, they act as precursor molecules that can release an antimicrobial peptide after proteolytic cleavage. But their structural features clearly distinguish them from defensins. The discovery of cathelicidins commenced after the isolation of the antimicrobial peptide Bac5 from bovine neutrophils, and the realization that they are cleaved from inactive precursors (Zanetti *et al.* 1990; Frank *et al.* 1990). In humans, the cathelicidin family is limited to a single gene, CAMP, with the protein product most frequently referred to as hCAP18/LL-37 (Cowland *et al.* 1995). The expression of hCAP18/LL-37 was initially detected directly in human skin keratinocytes only at sites of inflammation, although more recently it has also been found to be constitutively produced in specialized keratinocytes such as those of the nail or in neonatal skin and in eccrine glands (Dorschner *et al.* 2004). The first mature peptide isolated from this gene product was referred to as LL-37 because it begins with two leucine residues and is 37 amino acids long. This peptide forms a linear amphipathic α -helix. The primary structure of LL-37, LLGDF-FRKS-K-EKIGK-EFKRI-VQRIK-DFLRN-LVPRT-ES, was given along with other α -helical cathelicidin-derived antimicrobial peptides. The cathelicidin proteins are characterized by a highly conserved N-terminal domain of about 100 amino acid residues. This 14 kDa cathelin-like domain is flanked by a signal peptide domain (approximately 30 residues long) on its N-terminus, and by an antimicrobial peptide region on its C-terminus. This signal peptide is cleaved off once it has fulfilled its purpose of targeting the cathelicidin to storage granules or to the exterior of the cell. The formation of two disulfide-bridges finalizes post-transcriptional processing. Cathelicidin proteins have been extensively studied in mammals to date. However, cathelicidins were also recently found in chicken and three species of fish (Zanetti *et al.* 1991).

Defensins were first isolated from seeds of monocot and dicot species (Terras *et al.* 1995). Typical plant defensins form two additional disulfide bonds. Plant and arthropod defensins consist of a $\beta\alpha\beta\beta$ pattern, whereas mammalian β -defensins contain an N-terminal α -helix and an overall $\alpha\beta\beta\beta$ -fold. Plant defensins are small 45–54 amino acids long cationic peptides. Defensins are widely distributed among dicots and monocots. The genome of *A. thaliana*

alone was shown to encode more than 300 defensin-like (DEFL) peptides, 78% of which have a CS $\alpha\beta$ motif (Silverstein *et al.* 2005). Unlike animal defensins, few plant defensins are active against bacteria (Franco *et al.* 2006). Instead, the most common activity of these peptides is directed against diverse fungi (Osborn *et al.* 1995). Besides these antimicrobial activities, specific defensins have been reported to inhibit protein synthesis (Mendez *et al.* 1990). Plant defensins also influence plant growth and development. Altered expression of the tomato defensin *DEF2* in transgenic tomato plants reduced pollen viability and seed production (Stotz *et al.* 2009). Constitutive expression of this defensin had pleiotropic effects on plant development. In contrast to the above examples, the following developmental process has been studied at the mechanistic level: S-locus Cys-rich (SCR) protein as DEFL peptide for interaction with SPK, S-locus receptor kinase (SRK) to trigger self-incompatibility, a response that prevents self-fertilization via inhibition of pollen tube growth and that favors outcrossing (Nasrallah 2002). Also, L3 loop connecting β 2 and β 3 and the α -helical region are important for recognition of SCR by SRK (Sato *et al.* 2004). PR-13 is closely related to the defensin peptide NaD1 and more distantly related to thionins. Silencing of the tobacco defensin PR-13 resulted in increased susceptibility of *Nicotiana attenuata* to *Pseudomonas syringae* pv. tomato DC3000 under glasshouse conditions and increased susceptibility to opportunistic *Pseudomonas* spp. and mortality in the native habitat (Rayapuram *et al.* 2008). Based on these data, defensins are clearly important for plant defense, and their evolutionary diversification is the basis for various ecological functions. Defensins were originally grouped with the thionins and defined as g-thionins. However, structural comparisons to insect defensins required a reclassification of this group of AMPs as defensins (Bruix *et al.* 1993).

The first AMP isolated from plants was a thionin from the endosperm of wheat. Thionins have homologues from various taxa, such as viscotoxins, phoratoxins, and crambins (Mendez *et al.* 1990; Bruix *et al.* 1993). Thionins have four disulfide bonds from cereals and *Pyrularia pubera*, while other dicotyledonous thionins have three disulfide bonds. Thionins from the endosperm and leaf of barley inhibit the fungi *Drechslera teres*, a pathogen of barley, and *Thielaviopsis*

paradoxa, a pathogen of sugar cane (Bohlmann *et al.* 1988). Thionins also affect organisms other than plant pathogens. Purothionins are toxic to small mammals when injected intravenously or intraperitoneally but not when administered orally (Rayapuram *et al.* 2008). The mature thionin peptides are generally 45-47 amino acids in length and derived from preproproteins that contain an N-terminal signal peptide for targeting to the endoplasmic reticulum (ER) and a C-terminal acidic peptide that is thought to neutralize the activity of the cationic thionin. Also, thionins

are targeted to the vacuole to the action of basic chitinase and β -1,3-glucanase (Romero *et al.* 1997).

Transgenic plants expressing antimicrobial peptides

Antimicrobial peptides (AMPs) have been the object of attention in past years as candidates for plant protection products. Thus, gene constructions including sequences coding for AMPs have been expressed on model or crop plants providing different degrees of protection against plant pathogens. Transgenic plants introduced with defensin

Table 2. Summary of genetic engineering for antimicrobial peptides (AMPs) from plants.

AMP/ Signal sequence	Source for defensin	Transgenic plant	Promoter	Pathogens/pests tested*	Reference
BrD1	<i>Brassica rapa</i>	Rice	Rice cytochrome C	<i>Nilaparvata lugens</i> (brown planthopper insect)	Choi <i>et al.</i> 2009
RsAFP2	Radish	Wheat/rice	Maize ubiquitin/ CaMV 35S	<i>Fusarium graminearum</i> , <i>Rhizoctonia cerealis</i> , <i>Magnaporthe oryzae</i> , <i>Rhizoctonia solani</i> , <i>Alternaria longipes</i>	Li <i>et al.</i> 2011 Jha and Chattoo 2010 Terras <i>et al.</i> 1995
Dm-AMP1	<i>Dahlia merckii</i>	Rice/papaya	Maize ubiquitin/ D35S	<i>Magnaporthe oryzae</i> , <i>Rhizoctonia solani</i> , <i>Phytophthora palmivora</i>	Jha <i>et al.</i> 2009 Zhu <i>et al.</i> 2007
MsDef1	Alfalfa	Tomato	CaMV 35S	<i>Fusarium oxysporum</i>	Abdallah <i>et al.</i> 2010
NmDef02	<i>Nicotiana megalosiphon</i>	Tobacco/potato	CaMV 35S	<i>Phytophthora parasitica</i> , <i>Peronospora hyoscyami</i> , <i>Phytophthora infestans</i> , <i>Alternaria solani</i>	Portieles <i>et al.</i> 2010
WjAMP-1	Wasabi	Melon	CaMV 35S	<i>Fusarium oxysporum</i> , <i>Alternaria solani</i>	Ntui <i>et al.</i> 2010
cdef1	Chili	Tomato	CaMV 35S	<i>Fusarium sp.</i> , <i>Phytophthora infestans</i>	Zainal <i>et al.</i> 2009
BjD	Mustard	Peanut	CaMV 35S	<i>Phaeoisariopsis personata</i> , <i>Cercospora arachidicola</i> ,	Swathi Anuradha <i>et al.</i> 2008
alfAFP	Alfalfa	Potato	FMV35S	<i>Verticillium dahliae</i>	Gao <i>et al.</i> 2000
DRR206	Pea	Canola	CaMV 35S	slight increase in resistance to <i>Leptosphaeria maculans</i>	Wang <i>et al.</i> 1999
wasabi defensin	Wasabi	Rice	Ubiquitin-1	<i>Magnaporthe grisea</i>	Kanzaki <i>et al.</i> 2002
MiAMP1	<i>Macadamia integrifolia</i>	Canola	E12 Ω	<i>Leptosphaeria maculans</i>	Kazan <i>et al.</i> 2002

Note: CaMV = Cauliflower Mosaic Virus, E12 Ω = the 5' enhancer sequence from CaMV35S promoter + omega sequence from TMV, D35S = double CaMV35S + omega sequence from TMV, FMV = Figwort Mosaic Virus, ND = Not Determined, TSP=total soluble protein, * the transgenic plant displayed increased resistance to the tested organisms unless otherwise specified.

Table 3. Summary of genetic engineering for antimicrobial peptides (AMPs) from animals.

AMP	Source for AMP	Transgenic plant	Signal sequence	Promoter	Pathogens/pests tested*	Reference
Tachyplesin I	Horseshoe crabs	Potato	Barley α -hordothionin	CaMV 35S	Slight increase in resistance to <i>Erwinia</i> spp.	Allefs <i>et al.</i> 1996
Cecropin A	Giant silk moth	Tobacco	Cecropin B	CaMV 35S	No significant increase in resistance to <i>Pseudomonas syringae</i> pv. <i>tabaci</i>	Hightower <i>et al.</i> 1994
MB39	Synthetic	Tobacco	Barley α -amylase	PiII	<i>Pseudomonas syringae</i> pv. <i>tabaci</i>	Huang <i>et al.</i> 1997
Cecropin B	Giant silk moth	Tobacco/ Rice	Barley leaf thionin/ rice chitinase	CaMV 35S/ E7QIn	No significant increase in resistance to <i>Pseudomonas syringae</i> and <i>Pseudomonas solanacearum</i> / <i>Xanthomonas oryzae</i> pv. <i>oryzae</i>	Florack <i>et al.</i> 1995 / Sharma <i>et al.</i> 2000
Sarcotoxin IA	Flesh fly	Tobacco	Tobacco PR protein	E12Q	<i>Pseudomonas syringae</i> pv. <i>tabaci</i> and <i>Erwinia carotovora</i>	Ohshima <i>et al.</i> 1999
Attacin E	Giant silk moth	Pear	Attacin E	2 x CaMV35S	<i>Erwinia amylovora</i>	Reynold <i>et al.</i> 1999
D4E1	Synthetic	Tobacco	not specified	D35S	<i>Aspergillus flavus</i> and <i>Verticillium dahliae</i>	Cary <i>et al.</i> 2000
MB39	Synthetic	Apple	barley α -amylase	OSMp	<i>Erwinia amylovora</i>	Liu <i>et al.</i> 2001
MSI-99	Synthetic analog of frog magainin 2	Banana / Tomato/ Tobacco	Pea vicilin / expressed in chloroplast	<i>Arabidopsis</i> ubq 3 / CaMV35S/ 16S rRNA promoter	<i>Fusarium oxysporum</i> and <i>Mycosphaerella musicola</i> / <i>Pseudomonas syringae</i> pv. <i>tomato</i> but not to <i>Phytophthora infestans</i> or <i>Alternaria solani</i> / <i>P. syringae</i> pv. <i>tabaci</i> ATCC	Chakrabarti <i>et al.</i> 2003 / Alan <i>et al.</i> 2004/ DeGray <i>et al.</i> 2001
Gallerimycin	Greater wax moth <i>Galleria mellonella</i>	Tobacco	Gallerimycin	inducible mannopine synthase promoter	<i>Erysiphe cichoracearum</i> and <i>Sclerotinia minor</i>	Langen <i>et al.</i> 2006
Heliomicin	<i>Heliothis virescens</i>	Tobacco	Tobacco PR1a	D35S	Slight resistance to <i>Cercospora nicotianae</i>	Banzet <i>et al.</i> 2002
Drosomycin	<i>Drosophila melanogaster</i>	Tobacco	Tobacco PR1a	D35S	Slight resistance to <i>Cercospora nicotianae</i>	Banzet <i>et al.</i> 2002
Human β -defensin 2	Human	<i>A. thaliana</i>	plant defensin -DmAMP1	CaMV35S	<i>Botrytis cinerea</i>	Aerts <i>et al.</i> 2007
hCAP18 LL-37	Human	<i>Brassica rapa</i> / Tomato	Role gene encoding cytokinin synthesis	CaMV35S	<i>P. carotovorum</i> subsp. <i>Carotovorum</i> / <i>Fusarium oxysporum</i> f. sp. <i>Lycopersici</i> / <i>Colletotrichum higginsianum</i> / <i>Rhizoctonia solani</i> / <i>X. campestris</i> pv. <i>vesicatoria</i>	Jung <i>et al.</i> 2012/ Jung 2013
Rabbit α -defensin	Rabbit	Tobacco	not reported	CaMV35S	Slight resistance to <i>Ralstonia solanacearum</i>	Fu <i>et al.</i> 1998

Note: CaMV = Cauliflower Mosaic Virus, E12Q = the 5' enhancer sequence from CaMV35S promoter + omega sequence from TMV, E7QIn = synthesized high expression vector, D35S = double CaMV35S + omega sequence from TMV, FMV = Figwort Mosaic Virus, PiII = promoter from proteinase II inhibitor gene, OSMp = osmotin promoter, ND = Not Determined, TSP=total soluble protein, * the transgenic plant displayed increased resistance to the tested organisms unless otherwise specified.

genes have been used to control *Alternaria longipes* (Terras *et al.* 1995), *V. dahliae* (Gao *et al.* 2000), *Heterobasidium annosum* (Elfstrand *et al.* 2001), *Leptosphaeria maculans* (Wang *et al.* 1999), *Phytophthora parasitica* (Park *et al.* 2002), *M. grisea* (Kanzaki *et al.* 2002), *Botrytis cinerea* and *Verticillium albo-atrum* (Turrini *et al.* 2004), *Alternaria solani* (Schaefer *et al.* 2005) (Table 2). The hevein *Pn-AMP* expressed in tobacco protects against *Phytophthora parasitica* (Koo *et al.* 2002), and barley hordothionin expression in tobacco confers protection against *C. michiganensis* and *Pseudomonas syringae* pv. *tabaci* (Carmona *et al.* 1993). Also, human cathelicidin antimicrobial peptide carried out the substitution Met37Leu in Chinese cabbage and tomato. The expression of this peptide in cabbage and tomato plants significantly inhibited the growth of *Pectobacterium carotovorum* subsp. *carotovorum* on the plant leaves, and it conferred resistance to several fungal pathogens (Jung *et al.* 2012; Jung 2013a, b) (Table 2, 3, Fig. 2A, 2B).

Since pathogens have the ability to overcome gene-for-

gene host defense mechanisms in the field by undergoing mutations in the cognate avirulence genes, any transgenic model based on such resistance-conferring genes may easily be evaded by pathogens. On the other hand, this problem is less likely to occur in transgenic plants overexpressing genes with more general antimicrobial activity. Antimicrobial peptides from microbial origin expressed in plants are limited.

Concluding and Future Prospects

AMPs could play strong roles in agriculture as plant protection agents. In response to a microbial attack, plants activate a complex series of responses that lead to the local and systemic induction of a broad spectrum of antimicrobial defenses (Montesinos 2007). When induced defense responses are rapidly and coordinately triggered during a given plant-pathogen interaction, plants become broadly resistant to diseases. These defense responses include the strengthening

Fig. 2. (A) Temporal development of soft rot in Chinese cabbage (“Osome”) plants when the leaves were inoculated with *Pectobacterium carotovorum* subsp. *carotovorum*. B21–B24 Transgenic homozygous lines. (B) Disease development in transgenic homozygous lines containing the LL-37 gene, which were inoculated with 10^4 , 10^6 , or 10^8 CFU/mL of *P. carotovorum* subsp. *carotovorum*. Disease manifestations were scored 3 days after inoculation. RG water, WT wild-type plant, B21–B24 LL-37 transgenic homozygous lines. Disease lesions: 0, no lesion; 1, lesion size 0.1–0.5 cm; 2, 0.5–1.5 cm; 3, 1.5–3.5 cm; 4, 3.5–5.5 cm; 5, 5.5–8.5 cm; and 6, over 8.5 cm or plant dead. (Jung *et al.* 2012).

of mechanical barriers, oxidative burst, and production of antimicrobial compounds (Park 2005). Some research have been performed to bolster plant defenses against bacteria and fungi by genetically engineering plants to express antimicrobial peptides (Lee *et al.* 2008). Until now, studies involving the enhancement of resistance to various bacterial, fungal, and oomycete pathogens by the expression of antimicrobial peptides have been reported for rice, tobacco, poinsettia, banana, Chinese cabbage, tomato and more host species (Chakrabarti *et al.* 2003; Liang *et al.* 2002; Jung *et al.* 2012; Jung 2013b). Indeed these transgenic plants demonstrated reliable immunity to corresponding pathogen. However, exploitation of the great number of AMPs as active ingredients of pesticides has not been fully explored yet. Although majority of AMPs with potential uses have been studied at the *in vitro* level, fewer compounds have been tested on plant pathosystems, and only a few are in the market. Needless to say, more research are needed to address these areas. Further research area to explore with AMPs are its potential application for pharmaceutical purposes, and development of compounds suitable for agricultural use as pesticide ingredients. AMPs can be developed and processed commercially as less toxic and more stable compounds by improving preparative synthesis and biotechnological procedures using microbial systems or transgenic crops as plant factories.

ACKNOWLEDGMENT

This work was supported by a grant from the Next-Generation BioGreen 21 Program (No. PJ008085), Rural Development Administration, Republic of Korea.

REFERENCES

- Abdallah NA, Shah D, Abbas D, Madkour M. 2010. Stable integration and expression of a plant defensin in tomato confers resistance to fusarium wilt. *GM crops* 1: 344-350.
- Aerts AM, Thevissen K, Bresseleers SM, Sels J, Wouters P, Cammue BPA, François IEJA. 2007. *Arabidopsis thaliana* plants expressing human-defensin-2 are more resistant to fungal attack: functional homology between plant and human defensins. *Plant Cell Reports* 26: 1391-1398.
- Agerberth B, JY Lee, T Bergman, M Carlquist, HG Boman, V Mutt, H Jornvall. 1991. Amino acid sequence of PR-39. Isolation from pig intestine of a new member of the family of proline-arginine-rich antibacterial peptides. *Eur. J. Biochem.* 202: 849-854.
- Agrios GN. 2005. *Plant Pathology*. Academic Press, London.
- Alan AR, Blowers A, Earle ED. 2004. Expression of a magainin-type antimicrobial peptide gene (MSI-99) in tomato enhances resistance to bacterial speck disease. *Plant Cell Reports* 22: 388-396.
- Allefs S, De Jong E, Florack E, Hoogendoorn C, Stiekema W. 1996. *Erwinia* soft rot resistance of potato cultivars expressing antimicrobial peptide tachyplesin I. *Mol. r Breed.* 2: 97-105.
- Arnold RR, Cole MF, McGhee JR. 1977. A bactericidal effect for human lactoferrin. *Science* 197: 263-265.
- Bahar AA, Ren D. 2013. Antimicrobial peptides. *Pharmaceuticals* 6: 1543-1575.
- Balls AK. 1942. A crystalline protein obtained from a lipoprotein of wheat flour. *Cereal Chem.* 19: 279-288.
- Banzet N, Latorse MP, Bulet P, Francois E, Derpierre C, Dubald M. 2002. Expression of insect cystein-rich antifungal peptides in transgenic tobacco enhances resistance to a fungal disease. *Plant Science* 162: 995-1006.
- Bohlmann H, Clausen S, Behnke S, Giese H, Hiller C, Reimann-Philipp U, Schrader G, Barkholt V, Apel K. 1988. Leaf-specific thionins of barley-a novel class of cell wall proteins toxic to plant-pathogenic fungi and possibly involved in the defence mechanism of plants. *EMBO J.* 7: 1559-1565.
- Brouwer CP, Rahman M, Welling MM. 2011. Discovery and development of a synthetic peptide derived from lactoferrin for clinical use. *Peptides* 32: 1953-1963.
- Bruix M, Jimenez MA, Santoro J, Gonzalez C, Colilla FJ, Mendez E, Rico M. 1993. Solution structure of gamma 1-H and gamma 1-P thionins from barley and wheat endosperm determined by 1H-NMR: a structural motif common to toxic arthropod proteins. *Biochemistry* 32: 715-724.
- Carmona MJ, Molina A, Fernández JA, López-Fando JJ, García-Olmedo F. 1993. Expression of the alpha-thionin gene from barley in tobacco confers enhanced resistance

- to bacterial pathogens. *Plant J.* 3: 3-457.
- Cary JW, Rajasekaran K, Jaynes JM, Cleveland TE. 2000. Transgenic expression of a gene encoding a synthetic antimicrobial peptide results in inhibition of fungal growth *in vitro* and *in planta*. *Plant Science* 154: 171-181.
- Chakrabarti A, Ganapathi TR, Mukherjee PK, Bapat VA. 2003. MSI-99, a magainin analogue, imparts enhanced disease resistance in transgenic tobacco and banana. *Planta* 216: 587-596.
- Choi MS, Kim YH, Park HM, Seo BY, Jung JK, Kim ST, Kim MC, Shin DB, Yun HT, Choi IS, Kim CK, Lee JY. 2009. Expression of BrD1, a plant defensin from *Brassica rapa*, confers resistance against brown planthopper (*Nilaparvata lugens*) in transgenic rices. *Mol. Cells* 28: 131-137.
- Clark DP, S Durell, WL Maloy, M Zasloff. 1994. Ranalexin. A novel antimicrobial peptide from bullfrog (*Rana catesbeiana*) skin, structurally related to the bacterial antibiotic, polymyxin. *J. Biol. Chem.* 269: 10849-10855.
- Conlon JM, Sonnevend A. 2010. Antimicrobial peptides in frog skin secretions. *Methods Mol. Biol.* 618: 3-14.
- Conlon JM, T Halverson, J Dulka, JE Platz, FC Knoop. 1999. Peptides with antimicrobial activity of the brevinin-1 family isolated from skin secretions of the southern leopard frog, *Rana sphenoccephala*. *J. Pept. Res.* 54: 522-527.
- Cowland JB, Johnsen AH, Borregaard N. 1995. hCAP-18, a cathelin/ pro-bactenecin-like protein of human neutrophil specific granules. *FEBS Lett.* 368: 173-6.
- DeGray G, Rajasekaran K, Smith F, Sanford J, Daniell H. 2001. Expression of an antimicrobial peptide via the chloroplast genome to control phytopathogenic bacteria and fungi. *Plant Physiol.* 127: 852-862.
- Dorschner RA, Lopez-Garcia B, Massie J, Kim C, Gallo RL. 2004. Innate immune defense of the nail unit by antimicrobial peptides. *J. Am. Acad. Dermatol.* 50: 343-8.
- Dubos RJ. 1939. Studies on a bactericidal agent extracted from a soil bacillus: II. Protective effect of the bactericidal agent against experimental *Pneumococcus* infections in mice. *J. Exp. Med.* 70: 11-17.
- Dubos RJ, Hotchkiss RD. 1941. The production of bactericidal substances by aerobic sporulating bacilli. *J. Exp. Med.* 73: 629-640.
- Elfstrand M, Fossdal C, Swedjemark G, Sitbon F, Clapham D, Olsson O, Sharma P, Lönneborg A, von Arnold S. 2001. Identification of candidate genes for use in molecular breeding – A case study with the Norway spruce defensin-like gene, spi 1. *Silvae Genet.* 50: 75-81.
- Fahrner RL, T Dieckmann, SS Harwig, RI Lehrer, D Eisenberg, J Feigon. 1996. Solution structure of protegrin-1, a broad-spectrum antimicrobial peptide from porcine leukocytes. *Chem. Biol.* 3: 543-550.
- Florack DE, Allefs S, Bollen R, Bosch D, Visser B, Stiekema WJ. 1995. Expression of giant silkworm cecropin B genes in tobacco. *Transgenic Research* 4: 132-141.
- Franco OL, Murad AM, Leite JR, Mendes PA, Prates MV, Bloch C Jr. 2006. Identification of a cowpea gamma-thionin with bactericidal activity. *FEBS J.* 273: 3489-3497.
- Frank RW, Gennaro R, Schneider K, Przybylski M, Romeo D. 1990. Amino acid sequences of two proline-rich bactericidins. *J. Biol. Chem.* 265: 18871-18874.
- Fu RZ, Peng YF, Cao GC, Chen CX, Ma JS, Zhang LM, Li WB, Sun YR. 1998. Expression of rabbit defensin NP-1 gene in transgenic tobacco plants and its activity against bacterial wilt. *Chin. Sci. Bull.* 43: 1519-1522.
- Gao A, Hakimi SM, Mittanck CA. 2000. Fungal pathogen protection in potato by expression of a plant defensin peptide. *Nat. Biotechnol.* 18: 1307-1310.
- Gesell J, M Zasloff, SJ Opella. 1997. Two-dimensional ¹H NMR experiments show that the 23-residue magainin antibiotic peptide is an alpha-helix in dodecylphosphocholine micelles, sodium dodecylsulfate micelles, and trifluoroethanol/water solution. *J. Biomol. NMR* 9: 127-135.
- Gordon YJ, Romanowski EG, McDermott AM. 2005. A review of antimicrobial peptides and their therapeutic potential as anti-infective drugs. *Curr. Eye Res.* 30: 505-515.
- Groves ML, Peterson RF, Kiddy CA. 1965. Polymorphism in the red protein isolated from milk of individual cows. *Nature* 207: 1007-1008.
- Gudmundsson GH, Agerberth B, Odeberg J, Bergman T, Olsson B, Salcedo R. 1996. The human gene FALL39 and processing of the cathelin precursor to the antibacterial peptide LL-37 in granulocytes. *Eur. J. Biochem.* 238: 325-332.
- Hancock RE. 1997. Peptide antibiotics. *Lancet* 349: 418-422.
- Hancock RE, Lehrer R. 1998. Cationic peptides: A new source of antibiotics. *Trends Biotechnol.* 16: 82-88.

- Hightower R, Baden C, Penzes E, Dunsmuir P. 1994. The expression of cecropin peptide in transgenic tobacco does not confer resistance to *Pseudomonas syringae* pv. *tabaci*. *Plant Cell Reports* 13: 295-299.
- Hirsch JG. 1956. Phagocytin: A bactericidal substance from polymorphonuclear leucocytes. *J. Exp. Med.* 103: 589-611.
- Hotchkiss RD, Dubos RJ. 1940. Fractionation of the bactericidal agent from cultures of a soil *Bacillus*. *J. Biol. Chem.* 132: 791-792.
- Holak TA, A Engstrom, PJ Kraulis, G Lindeberg, H Bennich, TA Jones, AM Gronenborn, GM Clore. 1988. The solution conformation of the antibacterial peptide cecropin A: a nuclear magnetic resonance and dynamical simulated annealing study. *Biochemistry* 27: 7620-7629.
- Huang Y, Nordeen RO, Di M, Owens LD, McBeath JH. 1997. Expression of an engineered cecropin gene cassette in transgenic tobacco plants confers disease resistance to *Pseudomonas syringae* pv. *tabaci*. *Phytopathology* 87: 494-499.
- Hwang PM, Zhou N, Shan X, Arrowsmith CH, Vogel HJ. 1998. Three-dimensional solution structure of lactoferricin B, An antimicrobial peptide derived from bovine lactoferrin. *Biochemistry* 37: 4288-4298.
- Kang SJ, Won HS, Choi WS, Lee BJ. 2009. De novo generation of antimicrobial LK peptides with a single tryptophan at the critical amphipathic interface. *J. Pept. Sci.* 15: 583-588.
- Kanzaki H, Nirasawa S, Saitoh H. 2002. Overexpression of the wasabi defensin gene confers enhanced resistance to blast fungus (*Magnaporthe grisea*) in transgenic rice. *Theor. Appl. Genet.* 105: 809-814.
- Kazan K, Rusu A, Marcus JP, Goulter KC, Manners JM. 2002. Enhanced quantitative resistance to *Laptosphaeria maculans* conferred by expression of a novel antimicrobial peptide in canola (*Brassica napus* L.). *Mol. Breeding* 10: 63-70.
- Koo JC, Chun HJ, Park HC. 2002. Over-expression of a seed specific hevein-like antimicrobial peptide from *Pharbitis nil* enhances resistance to a fungal pathogen in transgenic tobacco plants. *Plant Mol. Biol.* 50: 441-452.
- Jha S, Chattoo BB. 2010. Expression of a plant defensin in rice confers resistance to fungal phytopathogens. *Transgenic Res.* 19: 373-384.
- Jha S, Tank HG, Prasad BD, Chattoo BB. 2009. Expression of Dm-AMP1 in rice confers resistance to *Magnaporthe oryzae* and *Rhizoctonia solani*. *Transgenic Res.* 18: 59-69.
- Jung YJ. 2013a. Transcriptional profiling of soft-rot resistant transgenic chinese cabbage (*Brassica rapa* L.) constitutively over expressing a human cathelicidin antimicrobial peptide (hCAP18/ LL-37). *Plant Breed. Biotech.* 1: 80-90.
- Jung YJ. 2013b. Enhanced resistance to bacterial pathogen in transgenic tomato plants expressing cathelicidin antimicrobial peptide. *Biotech. Bioproc. Eng.* 18: 616-625.
- Jung YJ, Lee SY, Moon YS, Kang KK. 2012. Enhanced resistance to bacterial and fungal pathogens by overexpression of a human cathelicidin antimicrobial peptide (hCAP18/LL-37) in Chinese cabbage. *Plant Biotechnol. Rep.* 6: 39-46.
- Langen G, Imani J, Altincicek B, Kieseritzky G, Kogel K, Vilcinskis A. 2006. Transgenic expression of gallerimycin, a novel antifungal insect defensin from the greater wax moth *Galleria mellonella*, confers resistance to pathogenic fungi in tobacco. *Biol. Chem.* 387: 549-557.
- Lee SC, Hwang IS, Choi HW, Hwang BK. 2008. Involvement of the pepper antimicrobial protein CaAMP1 gene in broad spectrum disease resistance. *Plant Physiol.* 143: 1004-1020.
- Li Z, Zhou M, Zhang Z, Ren L, Du L, Zhang B, Xu H, Xin Z. 2011. Expression of a radish defensin in transgenic wheat confers increased resistance to *Fusarium graminearum* and *Rhizoctonia cerealis*. *Funct. Integr Genomics* 11: 63-70.
- Liang H, Catranis CM, Maynard CE, Powell WA. 2002. Enhanced resistance to the poplar fungal pathogen, *Septoria musiva*, in hybrid poplar clones transformed with genes encoding antimicrobial peptides. *Biotechnol. Lett.* 24: 383-389.
- Liu Q, Ingersoll J, Owens L, Salih S, Meng R, Hammerschlag F. 2001. Response of transgenic Royal Gala apple (*Malus X domestica* Borkh.) shoots carrying a modified cecropin MB39 gene, to *Erwinia amylovora*. *Plant Cell Reports* 20: 306-312.
- Lupetti A, Paulusma-Annema A, Welling MM, Dogterom-Ballering H, Brouwer CP, Senesi S, Van Dissel JT, Nibbering PH. 2003. Synergistic activity of the N-terminal peptide of human lactoferrin and fluconazole against *Candida* species. *Antimicrob. Agents Chemother.* 47: 262-267.

- Lupetti A, Paulusma-Annema A, Welling MM, Senesi S, van Dissel JT, Nibbering PH. 2000. Candidacidal activities of human lactoferrin peptides derived from the N terminus. *Antimicrob. Agents Chemother.* 44: 3257-3263.
- Maloy WL, Kari UP. 1995. Structure-activity studies on magainins and other host defense peptides. *Biopolymers.* 37: 105-122.
- Mendez E, Moreno A, Colilla F, Pelaez R, Limas GG, Mendez R, Soriano F, Salinas M, de Haro C. 1990. Primary structure and inhibition of protein synthesis in eukaryotic cell-free system of a novel thionin, gamma-thionin, from barley endosperm. *Eur. J. Biochem.* 194: 533-539.
- Montesinos E. 2007. Antimicrobial peptides and plant disease control. *FEMS Microbiol Lett* 270: 1-11.
- Nasrallah JB. 2002. Recognition and rejection of self in plant reproduction. *Science* 296: 305-308.
- Nguyen LT, Haney EF, Vogel HJ. 2011. The expanding scope of antimicrobial peptide structures and their modes of action. *Trends Biotechnol.* 29: 464-472.
- Nguyen LT, Schibli DJ, Vogel HJ. 2005. Structural studies and model membrane interactions of two peptides derived from bovine lactoferricin. *J. Pept. Sci.* 11: 379-389.
- Ntui VO, Thirukkumaran G, Azadi P, Khan RS, Nakamura I, Mii M. 2010. Stable integration and expression of wasabi defensin gene in "Egusi" melon (*Colocynthis citrullus* L.) confers resistance to *Fusarium* wilt and *Alternaria* leaf spot. *Plant Cell Reports* 29: 943-954.
- Ohshima M, Mitsuhara I, Okamoto M, Sawano S, Nishiyama K, Kaku H, Natori S, Ohashi Y. 1999. Enhanced resistance to bacterial diseases of transgenic tobacco plants overexpressing sarcotoxin IA, a bactericidal peptide of insect. *J. Biochem.* 125: 431-435.
- Ohtani S, Okada T, Yoshizumi H, Kagamiyama H. 1977. Complete primary structures of two subunits of purothionin a, a lethal protein for brewer's yeast from wheat flour. *J. Biochem.* 82: 753-767.
- Osborn RW, De Samblanx GW, Thevissen K, Goderis I, Torrekens S, Van Leuven F, Attenborough S, Rees SB, Broekaert WF. 1995. Isolation and characterisation of plant defensins from seeds of Asteraceae, Fabaceae, Hippocastanaceae and Saxifragaceae. *FEBS Lett.* 368: 257-262.
- Oyston PC, Fox MA, Richards SJ, Clark GC. 2009. Novel peptide therapeutics for treatment of infections. *J. Med. Microbiol.* 58: 977-987.
- Padovan L, Scocchi M, Tossi A. 2010. Structural aspects of plant antimicrobial peptides. *Curr. Protein Pept. Sci.* 11: 210-219.
- Pardi A, XL Zhang, ME Selsted, JJ Skalicky, PF Yip. 1992. NMR studies of defensin antimicrobial peptides. 2. Three-dimensional structures of rabbit NP-2 and human HNP-1. *Biochemistry* 31: 11357-11364.
- Park CH, Kang YH, Chun HJ. 2002. Characterization of a stamen-specific cDNA encoding a novel plant defensin in chinese cabbage. *Plant Mol. Biol.* 50: 59-69.
- Park JM. 2005. The hypersensitive response: a cell death during disease resistance. *Plant Pathol. J.* 21: 99-101.
- Park SH, Kim HE, Kim CM, Yun HJ, Choi EC, Lee BJ. 2002. Role of proline, Cysteine and a disulphide bridge in the structure and activity of the anti-microbial peptide gaegurin 5. *Biochem. J.* 368: 171-182.
- Portieles R, Ayra C, Gonzalez E, Gallo A, Rodriguez R, Chacón O, López Y, Rodriguez M, Castillo J, Pujol M, Enriquez G, Borroto C, Trujillo L, Thomma BP, Borrás-Hidalgo O. 2010. *NmDef02*, a novel antimicrobial gene isolated from *Nicotiana megalosiphon* confers high-level pathogen resistance under greenhouse and field conditions. *P. Biotech. J.* 8: 678-690.
- Ramamoorthy A, Thennarasu S, Lee DK, Tan A, Maloy L. 2006. Solid-state NMR investigation of the membrane-disrupting mechanism of antimicrobial peptides MSI-78 and MSI-594 derived from magainin 2 and melittin. *Biophys. J.* 91: 206-216.
- Rammelkamp CH, Weinstein L. 1942. Toxic effects of tyrothricin, gramicidin and tyrocidine. *J. Infect. Dis.* 71: 166-173.
- Rayapuram C, Wu J, Haas C, Baldwin IT. 2008. PR-13/Thionin but not PR-1 mediates bacterial resistance in *Nicotiana attenuata* in nature, and neither influences herbivore resistance. *Mol. Plant Microbe. Interact.* 21: 988-1000.
- Reynold JP, Mourgues F, Norelli J, Aldwinckle HS, Briset MN, Chevreau E. 1999. First evidence for improved resistance to fire blight in transgenic pear expression the attacin E gene from *Hyalophora cecropia*. *Plant Science* 149: 23-31.
- Romeo D, B Skerlavaj, M Bolognesi, R Gennaro. 1988. Structure and bactericidal activity of an antibiotic dodecapeptide purified from bovine neutrophils. *J. Biol. Chem.* 263: 9573-9575.
- Romero A, Alamillo JM, Garcia-Olmedo F. 1997. Processing

- of thionin precursors in barley leaves by a vacuolar proteinase. *Eur. J. Biochem.* 243: 202-208.
- Sato Y, Okamoto S, Nishio T. 2004. Diversification and alteration of recognition specificity of the pollen ligand SP11/SCR in self-incompatibility of Brassica and Raphanus. *Plant Cell* 16: 3230-3241.
- Schaefer SC, Gasic K, Cammue B, Broekaert W, van Damme EJM, Peumans WJ, Korban SS. 2005. Enhanced resistance to early blight in transgenic tomato lines expressing heterologous plant defense genes. *Planta* 222: 858-866.
- Scocchi M, D Romeo, M Zanetti. 1994. Molecular cloning of Bac7, a proline- and arginine-rich antimicrobial peptide from bovine neutrophils. *FEBS Lett.* 352: 197-200.
- Seo MD, Won HS, Kim JH, Mishig-Ochir T, Lee BJ. 2012. Antimicrobial Peptides for Therapeutic Applications: A Review. *Molecules* 17: 12276-12286.
- Selsted ME, MJ Novotny, WL Morris, YQ Tang, W Smith, JS Cullor. 1992. Indolicidin, a novel bactericidal tridecapeptide amide from neutrophils. *J. Biol. Chem.* 267: 4292-4295.
- Sharma A, Sharma R, Imamura M, Yamakawa M, Machii H. 2000. Transgenic expression of cecropin B, an antibacterial peptide from *Bombyx mori*, confers enhanced resistance to bacterial leaf blight in rice. *FEBS Letters* 484: 7-11.
- Silverstein KAT, Graham MA, Paape TD, VandenBosch KA. 2005. Genome organization of more than 300 defensin-like genes in Arabidopsis. *Plant Physiol.* 138: 600-610.
- Sipos D, M Andersson, A Ehrenberg. 1992. The structure of the mammalian antibacterial peptide cecropin P1 in solution, determined by proton-NMR. *Eur. J. Biochem.* 209: 163-169.
- Stec B, Markman O, Rao U, Heffron G, Henderson S, Vernon LP, Brumfeld V, Teeter MM. 2004. Proposal for molecular mechanism of thionins deduced from physico-chemical studies of plant toxins. *J. Pept. Res.* 64: 210-224.
- Stotz HU, Wang Y, Spence B. 2009. A defensin from tomato with dual function in defence and development. *Plant Mol. Biol.* 71: 131-143.
- Swathi Anuradha T, Divya K, Jami SK, Kirti PB. 2008. Transgenic tobacco and peanut plants expressing a mustard defensin show resistance to fungal pathogens. *Plant Cell Reports* 27: 1777-1786.
- Terras FRG, Eggermont K, Kovaleva V, Raikhel NV, Osborn RW, Kester A, Rees SB, Torrekens S, Van LF, Vanderleyden J, Cammue BPA, Broekaert WF. 1995. Small cysteine-rich antifungal proteins from radish: their role in host defense. *Plant Cell* 7: 573-588.
- Turrini A, Sbrana C, Pitto L. 2004. The antifungal Dm-AMP1 protein from *Dahlia merckii* expressed in *Solanum melongena* is released in root exudates and differentially affects pathogenic fungi and mycorrhizal symbiosis. *New Phytol.* 163: 393-403.
- Van Epps HL. 2006. Rene dubos: Unearthing antibiotics. *J. Exp. Med.* 203: 259.
- Wang Y, Nowak G, Culley D, Hadwiger LA, Fristensky B. 1999. Constitutive expression of pea defense gene DRR206 confers resistance to blackleg (*Leptosphaeria maculans*) disease in transgenic canola (*Brassica napus*). *Mol. Plant-Microbe Interact.* 12: 410-418.
- Yi GS, CB Park, SC Kim, C Cheong. 1996. Solution structure of an antimicrobial peptide buforin II. *FEBS Lett.* 398: 87-90.
- Zainal Z, Marouf E, Ismail I, Fei CK. 2009. Expression of the *Capsicum annum* (Chili) defensin gene in transgenic tomatoes confers enhanced resistance to fungal pathogens. *Am. J. P. Physiol.* 4: 70-79.
- Zanetti M. 2004. Cathelicidins, multifunctional peptides of the innate immunity. *J. Leukoc. Biol.* 75: 39-48.
- Zanetti M, Litteri L, Gennaro R, Horstmann H, Romeo D. 1990. Bactenecins, defense polypeptides of bovine neutrophils, are generated from precursor molecules stored in the large granules. *J. Cell Biol.* 111: 1363-1371.
- Zasloff M. 2002. Antimicrobial peptides of multicellular organisms. *Nature* 415: 389-395.
- Zeya HI, Spitznagel JK. 1963. Antibacterial and enzymatic basic proteins from leukocyte lysosomes: Separation and identification. *Science* 142: 085-1087.
- Zhao X, Wu H, Lu H, Li G, Huang Q. 2013. Lamp: A database linking antimicrobial peptides. *PLoS One* 8: e66557.
- Zhu YJ, Agbayani R, Moore PH. 2007. Ectopic expression of *Dahlia merckii* defensin DmAMP1 improves papaya resistance to *Phytophthora palmivora* by reducing pathogen vigor. *Planta* 226: 87-97.
- Zimmermann GR, P Legault, ME Selsted, A Pardi. 1995. Solution structure of bovine neutrophil beta-defensin-12: the peptide fold of the beta-defensins is identical to that of the classical defensins. *Biochemistry* 34: 13663-13671.